



Neutral α -mannosidase from fungus *Aspergillus fischeri* (NCIM 508) belongs to Class IIC subfamily

Shashidhara KS^{1,2*}, Prashanth Kumar HP¹, Sameer Srivastava², Gaikwad SM²

¹Department of Biotechnology, Saphthagiri College of Engineering, Bangalore, Karnataka, 560057 India.,

²Biochemical Sciences Division, CSIR-National Chemical Laboratory, Pune, Maharashtra, 411008 India

Abstract

BACKGROUND & OBJECTIVE: The α -mannosidase gene from *Aspergillus fischeri* was isolated by DNA and RNA method. **METHODOLOGY:** Cloning of gene, isolation of recombinant plasmid and sequencing of gDNA gene was carried out and bioinformatic analysis was done for gene to get intron-exon information and for sequence homology. **RESULTS:** The α -mannosidase gene from *Aspergillus fischeri* is 1990 base pair long and sequence analysis showed presence of three introns and four exons in the gene. BLAST analysis revealed 100% similarity to the *Neosartorya fischeri* (*A. fischeri*) and 94% with *Aspergillus fumigatus* class I α -mannosidase reported in NCBI Gene Data Bank. **CONCLUSION:** The neutral α -mannosidase from *Aspergillus fischeri* could belong to the Class IIC subfamily.

Keywords: α -mannosidase gene, *Aspergillus fischeri*, Cloning, Sequencing, gDNA.

@2012 BioMedAsia All right reserved

1. Introduction

Mannosidases are Glycohydrolases involved in the processing of mannose containing glycans *in vivo*. They are involved in the maturation and degradation of glycoprotein-linked oligosaccharides. Use of natural substrates for specificity determination, antibodies from sub cellular localization, specific enzyme inhibitors and mutant cell lines, has revealed the presence of different α -mannosidase isoenzymes. This suggests that α -mannosidases have multiple functions in glycoprotein metabolism¹. There are two classes of processing α -mannosidase class I and class II, based on their distinctive substrate specificity, responses to inhibitors, cat ion requirements, protein molecular weights, sub cellular localization, enzyme mechanisms and characteristic regions of conserved amino acid sequences^{1,2}. They belong to family 47 and family 38 glycosidase, respectively in Henrissat's glycosidase classification^{2,3}.

There is significant correlation of biochemical and physiological roles of the various member genes for each of these two classes. Processing mannosidases are located in both the ER and Golgi of mammalian cells. The differential activity of Class II α -mannosidase towards artificial substrates has important practical ramifications. Because Class II enzymes are readily assayed with aryl mannosides, they are almost always employed for the initial screening of new potential inhibitors. Class I enzymes are not as easily assayed, so good inhibitors of these may be missed or incompletely characterized¹.

The purification, characterization, chemical modification and tryptophan environment of α -mannosidase from *A. fischeri* has already been reported⁴⁻⁸. The enzyme is a Class II α -

mannosidase as revealed by the substrate specificity and independency on the metal ion/s for activity⁴. The present study describes the cloning of a high molecular weight α -mannosidase from *Aspergillus fischeri* (*Neosartorya fischeri*) and its characterization and proves that the enzyme belongs to Class IIC subfamily of α -mannosidase.

2. Material and Methods

2.1 Material

Ampicillin, Kanamycin, Tris, IPTG, X-gal, SDS, BSA, EDTA and Ethidium bromide were purchased from Sigma-Aldrich, USA. Agarose, restriction enzymes, T4 DNA ligase, RNase A and lysozyme were obtained from GIBCO-BRL (USA), Promega (USA) and Amersham (UK). Taq DNA polymerase was obtained from Bangalore Genei (India). Plasmid vectors pGEM-T Easy Vector is purchased from Stratagene (USA). All other chemicals and solvents of analytical grade were purchased from HIMEDIA, Qualigens Fine Chemicals and E-Merck Laboratories, India. Pipette tips and micro centrifuge tubes were purchased from Axygen (USA).

2.2 Isolation of genomic DNA

The fungal culture *Aspergillus fischeri* was grown in 500 ml of Basal medium (pH 5.0) at 28 °C for 7 days with continuous shaking at 180 rpm. DNA was isolated according to the method described by Hopwood *et al.*, 1985⁹ with slight modifications.

2.3 Total RNA Isolation

Total RNA from fungal mycelium was isolated using Trizol reagent. RNase free environment was created and maintained as described by Blumberg (1987)⁶. All glass and plasticware was DEPC (0.1% in water) treated overnight and autoclaved. The pestle and mortar were also DEPC treated and then baked at 300 °C for 6 h. All materials were dried in a vacuum oven.

2.3 cDNA first strand synthesis by Reverse Transcription

Complementary DNA (cDNA) is synthesized from a mature mRNA template in a reaction catalyzed by the enzyme reverse transcriptase. In the present study cDNA first strand was synthesized using ImProm-II™ Reverse Transcription System (Promega, USA). The reactions were set up as per the

*Corresponding author

Full Address :

Dr. Shashidhara, K.S,
Saphthagiri College of Engineering, Chikkasandra, Hesaraghatta Main Road, Bangalore, 560057, India
Phone: +91-80-28372801/2/3

E-mail: shashidaraks@saphthagiri.edu.in

manufacturer's guidelines.

2.4 Extraction and purification of DNA from agarose gels

The restriction digested DNA or PCR amplified products were run on an agarose gel in 1X TAE buffer. The gel was stained with ethidium bromide (0.5 µg ml⁻¹) and viewed using a hand held long wavelength UV illuminator. The fragment of interest were excised from the gel and weighed. A 50-200 mg gel slice was transferred to a 1.5 ml micro centrifuge tube and 0.5 ml GEX buffer (Auprep™ GEL^X kit, Life Technologies, USA) added and further processed to load DNA on column and eluted. The eluted DNA was stored at -20 °C.

2.5 Ligation reaction

The pGEM-T vector and PCR amplified product were mixed in a molar ratio of 1:2 and suspended in 5 µl of DNA ligase buffer. To the mixture, 1 unit of ligase was added and incubated for 4 h at 16 °C.

2.6 Escherichia coli transformation and selection

LB medium (50 ml) was inoculated with 1% of the overnight grown *E. coli* culture and allowed to grow till 0.5 O.D. at 600 nm. The competent *E. coli* cells are made and were transformed according to Sambrook *et al.*, 1989¹⁰.

In order to differentiate between the colonies containing recombinant plasmids from the non-recombinant colonies, blue white selection of the colonies was performed using ampicillin (40 µg/ml), 40 ml of a stock solution of X-gal (20 µg/ml in dimethylformamide) and 4 ml solution of isopropylthio-β-D-galactoside (IPTG) (200 µg/ml).

2.7 Isolation of plasmid DNA from E. coli cells

The alkaline lysis method of Sambrook *et al.*¹⁰ was improvised upon so that 12-24 samples could be processed conveniently for plasmid DNA extraction within 3 h, with yields of 5-30 µg per 1.5 ml culture depending on the host strain and the plasmid vector.

2.8 Restriction digestion of DNA

Plasmid DNA restriction digestion was set up as per recommendations of manufacturer.

2.9 Primer design for the α-mannosidase gene and its amplification

The whole genomic sequence of *Neosartorya fischeri* (*Aspergillus fischeri*) fungus is available in the NCBI website. The information on α-mannosidase gene reveals that it is Class I α-mannosidase. As reported in our earlier manuscripts⁶⁻⁸, it is proved that the α-mannosidase of molecular weight 69200 Da (on SDS PAGE) from *Aspergillus fischeri* is a Class II enzyme. To confirm whether this is the same enzyme as reported in genome sequence, the tryptic digest of enzyme and peptide mass fingerprinting was done with MALDI-TOF. The peptide masses so obtained were compared with the peptide masses of theoretical tryptic digest of the protein sequence of reported α-mannosidase. 11 peptide masses were exactly matched with each other. Again, the molecular mass of the enzyme (69200 Da) studied is same as the reported. Based on this information, the reported α-mannosidase in website was confirmed same as the enzyme under study and the gene (cDNA) was used for the primer design. The genomic DNA was used as a template for the amplification of the α-mannosidase gene of interest. The following primers were designed based on the cDNA sequence available from the NCBI Gene Bank database. The NCBI Gene ID, ACCESSION XM, VERSION XM, GI are 4594076.001267006, 001267006.1, 119500499 respectively. Primers with restriction site to be incorporated in the amplified product were flanked with restriction sites on the 5' end of the respective primers. The forward primer with Nde site (5' CAT ATG ATT CTA GGT CGC AGG CGC 3') and reverse primer with Hind III site (5'AAG CTT TGT ACG AGC AGA GGG ACG CTT 3') were used for the study.

2.10 Bioinformatic analysis

Nucleotide and amino acid sequence analysis was done using software pDRAW 32 and online bioinformatics analysis

facility available at www.justbio.com, www.expasy.org and www.ncbi.nlm.nih.gov.

3. Results and Discussion

3.1 Dissection of the α-mannosidase gene from A. fischeri

Genomic DNA was extracted and Primers were designed based on the gene sequence available in NCBI Gene data Bank as described in materials and methods. PCR reactions were set up with *A. fischeri* genomic DNA using all possible combinations of the forward and the reverse primers. A DNA fragment of ~2.0 Kb was amplified (Figure I) with

Forward primer

AfManF [5' ATG ATT CTA GGT CGC AGG CGC 3']

Reverse primer

AfManR-2 [5' TGT ACG AGC AGA GGG ACG C 3']

3.2 Cloning of α-mannosidase gene

The amplicon obtained by primers AfManF forward and AfManR-2 reverse was cloned in to pGEM-T Easy Vector (Promega, USA) by TA cloning and transformed in to *E. coli*. The fact that *Taq* polymerase adds up poly A tail to the amplicon has been exploited in TA cloning. The pGEM-T has poly T at the cloning sites which make it easy to bind to the poly A of the PCR amplified fragment with poly A tail. The positive clones were selected based on the blue-white selection using IPTG and X-Gal. The positive clones were then cross checked for the presence of the gene by isolating the plasmid and sequencing it (Figure II).

The nucleotide sequences of α-mannosidase genomic clones were searched for sequence similarity in NCBI Gene Bank database and they showed sequence similarity with reported *Neosartorya fischeri* (*Aspergillus fischeri*) (100%) and *Aspergillus fumigatus* (94%) Class I α-mannosidase fungal genes. Sequencing data analysis of gDNA of α-mannosidase showed that the clone was of 1990 bp long. The sequences upon alignment showed 100% sequence similarity with *Neosartorya fischeri* (*A. fischeri*) class I α-mannosidase. The calculated GC content (www.justbio.com) of the gene was 51.0 %. Manually, the intron / exon junctions in the clone were identified after doing BLAST search in gene bank. The α-mannosidase gDNA nucleotide sequences show presence of putative four exons and three introns. In the α-mannosidase

Figure I: PCR with *A. fischeri* genomic DNA template and primers AfManF and AfManR-2. DNA size marker (lane 1), PCR amplified product (lane 2)

gene clone the four exons spanned the nucleotide positions 4 - 719, 773 - 1566, 1633 - 1892 and 1942 - 1984. The putative introns spanned nucleotide positions 720 - 772, 1567 - 1632 and 1893 - 1941 (Suppl material S1).

Restriction analysis of α-mannosidase gDNA was done using bioinformatic software pDRAW32.

Analysis was limited to enzyme site cutting not more than two times in the sequence.

3.3 α-mannosidase gene isolation from A. fischeri by cDNA method

Total RNA was extracted from the *A. fischeri* and cDNA was synthesized as described in materials and methods. PCR reactions were set up with *A. fischeri* cDNA




Figure II: Nucleotide sequence of GDNA amplicon of α -mannosidase (1990 bp)

```

1 CATATGATTC TAGTCGCAGG CGTATCGCT ATGCTTACC AACCTTTATT
51 CTAATTCCTA TTACTCTTCT CTTTCTCTCG AGTCTACCGC CTGGCCGGGA
101 ACCTCCCTTT GTCGACGACC ACTCATACTA CGCAGAATAC CATCCCAAC
151 CCCCGCAGAA CGATAACCGT ATCCATTGGA CCAACATAC CGAAAGACAT
201 CCGTCCCCCA GTTTTATTCC TCTCCCTACT CCCGCGCCAT CCGCAATTCC
251 TCGTGCCAG TACGACTTCC CGAAGGAATC ATGGCTTGGC CGGAGAAAAA
301 GAGTGAACAG ACAGAAGGCC GTC AAGGAGG CCTTCACGCA TGCTGGGAAA
351 GGGTACAAGC AGCATGCTTG GATGCGCGAT GAACTTTCGC CCCTCAGTGC
401 TCGTTACAGA ACCACATTTG CTGGCTGGGC TGCCAATCTG GTGGACGCAC
451 TTGACACTCT TGTATTATG GGTATGGAAG ACGAGTTCAA GGATGCGTTG
501 CATGCCATTG AGAGCATTTG TTTTATACC ACCGATGCCA CCGAATGAA
551 TGATTTCGAG ACTACTATCC GATATGTAGG AGGGTTACTT GCGCTCACG
601 ACCTGACCGA CGGAATAAC CCGATCCTTC TGAAGAAGGC TGTGGAACTG
651 GCGGACGTGA TCTACGATGC TTTTGATACT AGCAATAGG ATGCCCCAGT
701 CTGCGTGGCA ATGGAGCAGT AAGCCCGCCA TTTGTTTCTG GAACCACTTA
751 AAGCTGCTGT TAACTAGGCC ATAGATCTGC GCGAGGCGCT AGCAATCAAC
801 CAAGCAGACA AACTATCCTC GCAGAATTGG GTTCATTGAA TCTAGAATTC
851 ACCAGGCTGT ACCAGCTGAC TCATGATCCC AAGTACTTTG ATGCGGTCAA
901 ACGAATTACA GACGTTCTCG ACGACGCTCA GAATAAGACC AAAATCCCTG
951 GGCTGTGGCC TATGATGGTC ACGCTCAGG ATTTAGAGTT CACAGCCCT
1001 CGCTTACTG TGGGTGGAAT GGCCGACTCA ACATATGAGT ATTTGCCGAA
1051 AGAGCATATG CTACTGGGAG CTCAGACGGA TCAATACCGC AAGATGTATG
1101 CCGCCGCGAT GGAGGCGATC AAGAGCGCT TATTGTTTCG ACCCATGATG
1151 AAAAACGGCG AGGATCTCCT CTTCCGCGGC AACACCCACA CGAGTCTCGC
1201 AACTAACGCG CCCCGGAGC CTCATGGA AATCTGAAA TGCTTTTTCG
1251 GGGGCACTGT GGGCATCGGG GCAAAGTTT TCAACCGCCC GGAGGAGCTC
1301 TCAATTGCAC GAAAGTAA CAGATGGCTG ATCTGGGCAT ACGATGTCAT
1351 GCCCACCGGA ATCATGCCG AATCATGCA TCTCAGCGCT TGCCAGAGTA
1401 TGGATAGTTG TGAATGGAG GAGCAGAAGT GGTACCAAGA TGTCGGCGCG
1451 CGACTGGCCA AAGGATCAA GGAGAGAGC ACTGTTCAAG AAGCAAAGC
1501 AATTATCCAG GAGAGTGGG TCAACTGG CTAATCTGAG ATATCGGATC
1551 CGAGATATT GCIGGGTAT GTATTGCT GACAATTGAT TTTCTGTTG
1601 CATCGGGTGT AGTAGTCTG ATATTGGTT TACAGCCGAG GCTATCGAGT
1651 CAGTTTCAAT CATGTACCGC ATCACTGGGC ACAAGAAATT GCAAGACGAT
1701 GCATGGAGAA TGTTTACAG CATTGAGAAA GCAACTGAAA CCAATCATC
1751 CCATGCCGCG ATAGACGAGC TCCGGGACGT GAAGACAACC CAGTAGATT
1801 ACATGGAGAG TTTCTGGCTG CAGAGACCT TGAAGTACTT CIATCTCGTC
1851 TTCTCCGAAC CAGGGCTTGT CAGCTTGGAT GATTACGTAC TGATGTGCT
1901 GTGTCTTGT GAGGCTCTG CCAAAGCTAA CGATTAAGCA GGAACACGGA
1951 AGCACATCCA TCAAGCGCT CCTCTGCTCG TACAAAGCTT

```

using all possible combinations of the forward and the reverse primers. A DNA fragment of ~1.8 Kb was amplified (Figure III) with Forward primer

AfManF [5'ATG ATT CTA GGT CGC AGG CGC 3']


Reverse primer

AfManR-2 [5' TGT ACG AGA GGG ACG C 3']

4. DISCUSSION

The α -mannosidase gene from *Aspergillus fischeri* was isolated

Figure III: PCR with *A. fischeri* cDNA template and primers AfManF and AfManR-2. DNA size marker (lane 1), PCR amplified product (lane 2).



by both genomic DNA and RNA method. Cloning of gene, isolation of recombinant plasmid and sequencing of genomic DNA gene showed that it is 1990 base pair long and its bioinformatic's analysis showed presence of three introns and four exons in the gene.

BLAST analysis revealed 100% similarity to the *Aspergillus fischeri* and 94% with *Aspergillus fumigatus* class I α -mannosidase reported in NCBI Gene Data Bank. But this sequence is not matching with any Class II α -mannosidase at gene level. Our previous studies have revealed that the enzyme is class II α -mannosidase.

So far, there are three subfamilies of genes known, Class IIA, IIB and IIC^{1,11,12}. Class II A α -mannosidase occurs only in higher eukaryotes,

involved in N-glycan synthesis in golgi and removes α , 1-3, 1-2, 1-6

linked mannose residues from N-glycans during their synthesis. Class II B α -mannosidase found only higher eukaryotes and involved in N-glycan breakdown, removal and recycling in the

cytoplasm, lysosome and vacuole. Class II C α -mannosidase found in higher and lower eukaryotes and involved in N-glycan breakdown, removal and recycling in the cytoplasm, lysosome and vacuole. They are heterogeneous enzymes with a diversity of functions and cellular location and show low sequence similarity with the other two groups of enzymes¹². Phylogenetic analysis of the sequences shows that the Class IIA and Class IIB subfamilies have diverged more recently than the Class IIC subfamily. A likely scenario is that a single common ancestor was duplicated after the divergence of lower eukaryotes, such as fungi, from the higher eukaryotes. The lower eukaryotes thus only contain the orthologue of the common ancestor¹³. As Class IIC found in higher and lower eukaryotes and has very low sequence similarity to the other two subfamilies and the present α -mannosidase from *Aspergillus fischeri* shows similar properties, hence the *Aspergillus fischeri* α -mannosidase could belong to the Class IIC subfamily.

Acknowledgement

Authors are thankful to Dr. BM Khan for providing lab facilities to carry out experiments during the course of this work. Financial support by CSIR (India) for SKS is duly acknowledged.

References

- Daniel PF, Winchester B, & Warren CD, Mammalian alpha-mannosidases--multiple forms but a common purpose?. *Glycobiol*, **4** (1994) 551.
- Moremen K, Trimble RB, & Herscovics A, Glycosidases of the asparagine-linked oligosaccharide processing pathway. *Glycobiol*, **4** (1994) 113.
- Henrissat BA, Updating the sequence-based classification of glycosyl hydrolases. *Biochem J*, **316** (1996) 695.
- Gaikwad SM, Keskar SS, & Khan MI, Purification and characterization of alpha-D-mannosidase from *Aspergillus* sp. *Biochim Biophys Acta*, **1250** (1995) 144.
- Gaikwad SM, Khan MI, & Keskar SS, Chemical modification of α -mannosidase from *Aspergillus* sp. *Biotechn. Appl Biochem*, **25** (1997) 105.
- Shashidhara KS, & Gaikwad SM, Fluorescence quenching and time-resolved fluorescence studies of alpha-mannosidase from *Aspergillus fischeri* (NCIM 508). *J Fluoresc*, **17** (2007) 599.
- Shashidhara KS, & Gaikwad SM, Class II alpha-mannosidase from *Aspergillus fischeri*: energetics of catalysis and inhibition. *Int J Biol Macromol*, **44** (2009) 112.
- Shashidhara KS, & Gaikwad SM, Conformational and functional transitions in class II alpha-mannosidase from *Aspergillus fischeri*. *J Fluoresc*, **20** (2010) 827.
- Hopwood DA, *Genetic manipulation of Streptomyces: a laboratory manual*. Norwich: John Innes Foundation, (1985).
- Sambrook J, Fritsch EF & Maniatis T, *Molecular cloning: a laboratory manual*, 2nd ed. Cold Spring Harbor, N.Y.: Cold Spring Harbor Laboratory Press, (1989).
- Eades CJ, & Hintz WE, Characterization of the class I alpha-mannosidase gene family in the filamentous fungus *Aspergillus nidulans*. *Gene*, **255**(1) (2000) 25.
- Eades CJ, Gilbert AM, Goodman CD, & Hintz WE, Identification and analysis of a class 2 alpha-mannosidase from *Aspergillus nidulans*. *Glycobiology*, **8**(1) (1998) 17.
- Eades CJ, & Hintz WE, Characterization of the α -mannosidase Gene Family in Filamentous Fungi: N-glycan Remodelling for the Development of Eukaryotic Expression Systems. *Biotechnol Bioprocess Eng*, **5** (2000) 227.